

LORAZEPAM (ATIVAN®)

| | |
|---------------------------------|--|
| THERAPEUTIC EFFECTS | Anticonvulsant |
| INDICATIONS | Major motor seizures when midazolam is not available. |
| CONTRAINDICATIONS | <ul style="list-style-type: none"> ✓ Hypersensitivity to lorazepam ✓ Acute narrow angle glaucoma ✓ Hypotension ✓ Respiratory failure <p>Relative Contraindications:</p> <ul style="list-style-type: none"> ✓ Respiratory distress ✓ Decreased LOC |
| PRECAUTIONS/SIDE EFFECTS | <ul style="list-style-type: none"> ✓ Respiratory depression ✓ Hypotension ✓ Sedation ✓ Nausea ✓ Vomiting |
| ADULT DOSAGE/ROUTE | Dilute 1:1 with an equal volume of NS and administer IV/IM* or 1-2 mg IO slowly and may repeat q 3-5 minutes, up to a maximum dose of 4 mg. |
| PEDIATRIC DOSAGE/ROUTE | Dilute 1:1 with an equal volume of NS and administer IV/IM*, PR, or 0.1 mg/kg IO q 3 minutes, up to a maximum dose of 2 mg. |

*IM lorazepam should not be diluted