

HDVCH EMS AND ED CARE FOR CHILD WITH CONGENITAL HEART DISEASE

- Cautious use of oxygen. Correlate with my normal saturation range. Titrate up slowly to maintain expected saturations.
- Contact medical control and consider 5mL/kg boluses if needed. Boluses can be repeated.
- Use Everbridge Page to notify team when call received from EMS.



GENERAL CARE FOR FAMILIES

- Wash your hands before touching my wound. Keep my wound clean and dry for 7 days after surgery. Do not submerge my wound under water for 4 weeks.
- Keep my chin off of my wound and cover with dry gauze until skin is completely healed.
- Notify the CHC if: I am drinking less than normal, saturations are lower than my normal range, my wound is red/has drainage, I have a fever above 101°F, if I have increased work of breathing or blueness, or with any other concerns.



EMERGENCY CARD FOR CHILD WITH CONGENITAL HEART DISEASE

IN AN EMERGENCY: CALL 911



MY CHD CHILD'S INFORMATION

Child's Name: _____

Fundamental Diagnosis: _____

Current Diagnosis: _____

Surgical Status:

Unrepaired Last Operated On: _____



MY SHUNT STATUS

I AM shunt dependent I AM NOT shunt dependent

MY NORMAL SATURATION RANGE

70% - 85% 80% - 90% Greater than 90%



ADDITIONAL CARE

I have a mechanical valve

INR Goal 2-3

INR Goal 2.5-4

Other INR goal: _____

Require SBE Prophylaxis for qualifying procedures.

(Single dose of Amoxicillin, unless allergy is present.

Child dose 50 mg/kg, Adult dose 2 grams).